

1993. ABSTRACT. Pages 16-17 in Symposium Contraception in Wildlife Management (October 26-28, 1993, Denver, Colorado)

EVALUATION OF RECOMBINANT BCG AS A LIVE VECTOR FOR ORAL IMMUNIZATION OF WHITE-TAILED DEER

BRAD E. JOHNS, LOWELL A. MILLER, and R. D. THOMPSON, U.S. Department of Agriculture, Animal and Plant Health Inspection Service, Animal Damage Control, Denver Wildlife Research Center, Denver, CO 80225
GARY J. KILLIAN, Pennsylvania State University, University Park, PA 16802

Abstract: A recombinant form of *Bacillus Calmette Guerin* (BCG), an attenuated bovine tuberculosis bacterium (*Mycobacterium bovis*), was used to test the feasibility of oral delivery of a vaccine to deer. The recombinant BCG expresses Osp A antigen, from a spirochete (*Borrelia burgdorferi*), on its surface as a model antigen and vector for a proposed vaccine that would deliver a genetically engineered immunocontraceptive to deer.

Ten white-tailed deer (*Odocoileus virginianus*) were tested and a class B containment facility used to ensure environmental safety. Deer were immunized as follows: 2 subcutaneously injected with Osp A (antigen controls); 2 subcutaneously injected with non-recombinant BCG (BCG controls); 2 subcutaneously injected with recombinant BCG containing Osp A; and 4 orally dosed with recombinant BCG containing Osp A. Osp A antigen control deer were given a booster dose 30 days after the prime dose; other treatment groups were administered a booster dose 90 days following the prime dose.

Injected Osp A in antigen controls proved to be highly immunogenic, producing high antibody titers. Recombinant BCG proved effective as a vaccine delivery vehicle, producing antibodies to Osp A by both subcutaneous injection and oral delivery. These Osp A antibody peak titers, measured by ELISA, were 1/1600 by injection and 1/6400 by oral delivery. Control deer that were housed with orally vaccinated deer demonstrated no BCG antibody response, indicating that recombinant BCG lacks infective properties. Acid-fast staining of fecal samples showed peak BCG shedding occurred at 24 hours and was 95% complete in 72 hours. While it appears feasible to use recombinant BCG as an oral vaccine vector, a DNA probe test would need to be developed to differentiate between bovine tuberculosis and genetically altered BCG before it could be used under free-ranging conditions.